


**XIth International Conference on
Elastic and Diffractive Scattering
Château de Blois, France, May 15 - 20, 2005**

EUCLIDEAN FORMULATION OF DIFFRACTIVE SCATTERING

E. MEGGIOLARO

*Dipartimento di Fisica, Università di Pisa, and I.N.F.N., Sezione di Pisa,
Largo Pontecorvo 3, I-56127 Pisa, Italy*



After a brief review (in the first part) of some relevant properties of the high-energy parton-parton scattering amplitudes, in the second part we shall discuss the infrared finiteness and some analyticity properties of the loop-loop scattering amplitudes in gauge theories, when going from Minkowskian to Euclidean theory, and we shall see how they can be related to the still unsolved problem of the s -dependence of the hadron-hadron total cross-sections.

1 Parton-parton scattering amplitudes

The parton-parton elastic scattering amplitude, at high squared energies s in the center of mass and small squared transferred momentum t (that is $s \rightarrow \infty$ and $|t| \ll s$, let us say $|t| \leq 1 \text{ GeV}^2$), can be described by the expectation value of two *infinite lightlike* Wilson lines, running along the classical trajectories of the colliding particles^{1,2,3}. However, this description is affected by infrared (IR) divergences, which are typical of $3 + 1$ dimensional gauge theories. One can regularize this IR problem by letting the Wilson lines coincide with the classical trajectories for partons with a non-zero mass m (so forming a certain *finite* hyperbolic angle χ in Minkowskian space-time: of course, $\chi \rightarrow \infty$ when $s \rightarrow \infty$), and, in addition, by considering *finite* Wilson lines, extending in proper time from $-T$ to T (and eventually letting $T \rightarrow +\infty$)^{4,5}. For example, the high-energy quark-quark elastic scattering amplitude $\mathcal{M}_{fi}$ is (explicitly indicating the colour indices i, j [initial] and i', j' [final] and the spin indices α, β [initial] and α', β' [final] of the colliding quarks):

$$\mathcal{M}_{fi} \underset{s \rightarrow \infty}{\sim} -i 2s \delta_{\alpha'\alpha} \delta_{\beta'\beta} g_M(\chi \rightarrow \infty; T \rightarrow \infty; t), \quad (1)$$

$$g_M(\chi; T; t) \equiv \frac{1}{[Z_W(T)]^2} \int d^2 \vec{z}_\perp e^{i\vec{q}_\perp \cdot \vec{z}_\perp} \langle [W_1^{(T)}(\vec{z}_\perp) - \mathbf{1}]_{i'i} [W_2^{(T)}(\vec{0}_\perp) - \mathbf{1}]_{j'j} \rangle, \quad (2)$$

where $t = -|\vec{q}_\perp|^2$, $\vec{q}_\perp$ being the tranferred momentum, and $\vec{z}_\perp = (z^2, z^3)$ is the distance between the two trajectories in the *transverse* plane (*impact parameter*). The two IR-regularized Wilson lines are defined as:

$$\begin{aligned} W_1^{(T)}(\vec{z}_\perp) &\equiv \mathcal{T} \exp \left[-ig \int_{-T}^{+T} A_\mu \left(z + \frac{p_1}{m} \tau \right) \frac{p_1^\mu}{m} d\tau \right], \\ W_2^{(T)}(\vec{0}_\perp) &\equiv \mathcal{T} \exp \left[-ig \int_{-T}^{+T} A_\mu \left(\frac{p_2}{m} \tau \right) \frac{p_2^\mu}{m} d\tau \right], \end{aligned} \quad (3)$$

where $\mathcal{T}$ stands for “time ordering” and $A_\mu = A_\mu^a T^a$; $z = (0, 0, \vec{z}_\perp)$; $p_1 = E(1, \beta, 0, 0)$ and $p_2 = E(1, -\beta, 0, 0)$ are the initial four-momenta of the two quarks [$s \equiv (p_1 + p_2)^2 = 2m^2(\cosh \chi + 1)$]. Finally, $Z_W(T)$ is a sort of Wilson-line’s renormalization constant:

$$Z_W(T) \equiv \frac{1}{N_c} \langle \text{Tr}[W_1^{(T)}(\vec{0}_\perp)] \rangle = \frac{1}{N_c} \langle \text{Tr}[W_2^{(T)}(\vec{0}_\perp)] \rangle. \quad (4)$$

The expectation values $\langle W_1 W_2 \rangle$, $\langle W_1 \rangle$, $\langle W_2 \rangle$ are averages in the sense of the QCD functional integrals:

$$\langle \mathcal{O}[A] \rangle = \frac{1}{Z} \int [dA] \det(Q[A]) e^{iS_A} \mathcal{O}[A], \quad (5)$$

where $Z = \int [dA] \det(Q[A]) e^{iS_A}$ and $Q[A]$ is the *quark matrix*.

The quantity $g_M(\chi; T; t)$ with $\chi > 0$ can be reconstructed from the corresponding Euclidean quantity $g_E(\theta; T; t)$, defined as a (properly normalized) correlation function of two (IR-regularized) Euclidean Wilson lines $\tilde{W}_1$ and $\tilde{W}_2$, i.e.,

$$\begin{aligned} g_E(\theta; T; t) &\equiv \frac{1}{[Z_{WE}(T)]^2} \int d^2 \vec{z}_\perp e^{i\vec{q}_\perp \cdot \vec{z}_\perp} \langle [\tilde{W}_1^{(T)}(\vec{z}_\perp) - \mathbf{1}]_{i'i} [\tilde{W}_2^{(T)}(\vec{0}_\perp) - \mathbf{1}]_{j'j} \rangle_E, \\ Z_{WE}(T) &\equiv \frac{1}{N_c} \langle \text{Tr}[\tilde{W}_1^{(T)}(\vec{0}_\perp)] \rangle_E = \frac{1}{N_c} \langle \text{Tr}[\tilde{W}_2^{(T)}(\vec{0}_\perp)] \rangle_E, \end{aligned} \quad (6)$$

where:

$$\begin{aligned} \langle \mathcal{O}[A^{(E)}] \rangle_E &= \frac{1}{Z^{(E)}} \int [dA^{(E)}] \det(Q^{(E)}[A^{(E)}]) e^{-S_A^{(E)}} \mathcal{O}[A^{(E)}], \\ Z^{(E)} &= \int [dA^{(E)}] \det(Q^{(E)}[A^{(E)}]) e^{-S_A^{(E)}}, \end{aligned} \quad (7)$$

$\theta \in]0, \pi[$ being the angle formed by the two trajectories in the Euclidean four-space, by an analytic continuation in the angular variables and in the IR cutoff^{5,6,7}:

$$\begin{aligned} g_E(\theta; T; t) &= g_M(\chi \rightarrow i\theta; T \rightarrow -iT; t), \\ g_M(\chi; T; t) &= g_E(\theta \rightarrow -i\chi; T \rightarrow iT; t). \end{aligned} \quad (8)$$

This result is derived under the assumption that the function g_M , as a function of the *complex* variable χ , is *analytic* in a domain $\mathcal{D}_M$ which includes the positive real axis ($\text{Re}\chi > 0, \text{Im}\chi = 0$) and the imaginary segment ($\text{Re}\chi = 0, 0 < \text{Im}\chi < \pi$); and, therefore, the function g_E , as a function of the *complex* variable θ , is *analytic* in a domain $\mathcal{D}_E = \{\theta \in \mathbf{C} \mid i\theta \in \mathcal{D}_M\}$, which includes the real segment ($0 < \text{Re}\theta < \pi, \text{Im}\theta = 0$) and the negative imaginary axis ($\text{Re}\theta = 0, \text{Im}\theta < 0$). The validity of this assumption is confirmed by explicit calculations in

perturbation theory⁶. Eq. (8) is then intended to be valid for every $\chi \in \mathcal{D}_M$ (i.e., for every $\theta \in \mathcal{D}_E$).

The above-reported relations allow to give a nice geometrical interpretation of the so-called *crossing symmetry*. Changing from a *quark* to an *antiquark* just corresponds, in our formalism, to substitute the corresponding Wilson line with its complex conjugate, i.e., to reverse the orientation of the Wilson line (and the colour indices):

$$[W_p^*(\vec{b}_\perp)]_{lk} = [W_p^\dagger(\vec{b}_\perp)]_{kl} = [W_{-p}(\vec{b}_\perp)]_{kl}. \quad (9)$$

Changing *quark* nr. 2 into an *antiquark* corresponds, in the Euclidean theory, to the substitution:

$$\theta \rightarrow \theta_2 = \pi - \theta, \quad (10)$$

and therefore, in the Minkowskian theory:

$$\chi \rightarrow \chi_2 = i\pi - \chi. \quad (11)$$

We thus find the following *crossing-symmetry* relation:

$$g_M^{(q\bar{q})}(\chi; T; t)_{i'l, lk} = g_M(i\pi - \chi; T; t)_{i'l, kl}. \quad (12)$$

[We must assume that the domain $\mathcal{D}_M$ also includes the half-line ($\text{Re}\chi < 0, \text{Im}\chi = \pi$).]

We close this section remarking that the *regularized* quantities $g_M(\chi; T; t)$ and $g_E(\theta; T; t)$, while being finite at any given value of T , are divergent in the limit $T \rightarrow \infty$. In some cases this IR divergence can be factorized out and one thus ends up with an IR-finite (physical) quantity.

2 Loop-loop scattering amplitudes

Differently from the parton-parton scattering amplitudes, which are known to be affected by infrared (IR) divergences, the elastic scattering amplitude of two colourless states in gauge theories, e.g., two $q\bar{q}$ meson states, is expected to be an IR-finite physical quantity. It was shown in Refs.^{8,9,10} that the high-energy meson-meson elastic scattering amplitude can be approximately reconstructed by first evaluating, in the eikonal approximation, the elastic scattering amplitude of two $q\bar{q}$ pairs (usually called “*dipoles*”), of given transverse sizes $\vec{R}_{1\perp}$ and $\vec{R}_{2\perp}$ respectively, and then averaging this amplitude over all possible values of $\vec{R}_{1\perp}$ and $\vec{R}_{2\perp}$ with two proper squared wave functions $|\psi_1(\vec{R}_{1\perp})|^2$ and $|\psi_2(\vec{R}_{2\perp})|^2$, describing the two interacting mesons. The high-energy elastic scattering amplitude of two *dipoles* is governed by the (properly normalized) correlation function of two Wilson loops $\mathcal{W}_1$ and $\mathcal{W}_2$, which follow the classical straight lines for quark (antiquark) trajectories:

$$\mathcal{M}_{(ll)}(s, t; \vec{R}_{1\perp}, \vec{R}_{2\perp}) \equiv -i \, 2s \int d^2\vec{z}_\perp e^{i\vec{q}_\perp \cdot \vec{z}_\perp} \left[\frac{\langle \mathcal{W}_1 \mathcal{W}_2 \rangle}{\langle \mathcal{W}_1 \rangle \langle \mathcal{W}_2 \rangle} - 1 \right], \quad (13)$$

where s and $t = -|\vec{q}_\perp|^2$ ($\vec{q}_\perp$ being the transferred momentum) are the usual Mandelstam variables. More explicitly the Wilson loops $\mathcal{W}_1$ and $\mathcal{W}_2$ are so defined:

$$\begin{aligned} \mathcal{W}_1^{(T)} &\equiv \frac{1}{N_c} \text{Tr} \left\{ \mathcal{P} \exp \left[-ig \oint_{\mathcal{C}_1} A_\mu(x) dx^\mu \right] \right\}, \\ \mathcal{W}_2^{(T)} &\equiv \frac{1}{N_c} \text{Tr} \left\{ \mathcal{P} \exp \left[-ig \oint_{\mathcal{C}_2} A_\mu(x) dx^\mu \right] \right\}, \end{aligned} \quad (14)$$

where $\mathcal{P}$ denotes the “*path ordering*” along the given path $\mathcal{C}$; $\mathcal{C}_1$ and $\mathcal{C}_2$ are two rectangular paths which follow the classical straight lines for quark [$X_{(+)}(\tau)$, forward in proper time τ] and

antiquark $[X_{(-)}(\tau)$, backward in τ] trajectories, i.e.,

$$\begin{aligned}\mathcal{C}_1 &\rightarrow X_{(\pm 1)}^\mu(\tau) = z^\mu + \frac{p_1^\mu}{m}\tau \pm \frac{R_1^\mu}{2}, \\ \mathcal{C}_2 &\rightarrow X_{(\pm 2)}^\mu(\tau) = \frac{p_2^\mu}{m}\tau \pm \frac{R_2^\mu}{2},\end{aligned}\tag{15}$$

and are closed by straight-line paths at proper times $\tau = \pm T$, where T plays the role of an IR cutoff, which must be removed at the end ($T \rightarrow \infty$). Here p_1 and p_2 are the four-momenta of the two quarks and of the two antiquarks with mass m , moving with speed β and $-\beta$ along, for example, the x^1 -direction:

$$\begin{aligned}p_1 &= m(\cosh \frac{\chi}{2}, \sinh \frac{\chi}{2}, 0, 0), \\ p_2 &= m(\cosh \frac{\chi}{2}, -\sinh \frac{\chi}{2}, 0, 0),\end{aligned}\tag{16}$$

where $\chi = 2 \operatorname{arctanh} \beta > 0$ is the hyperbolic angle between the two trajectories (+1) and (+2). Moreover, $R_1 = (0, 0, \vec{R}_{1\perp})$, $R_2 = (0, 0, \vec{R}_{2\perp})$ and $z = (0, 0, \vec{z}_\perp)$, where $\vec{z}_\perp = (z^2, z^3)$ is the impact-parameter distance between the two loops in the transverse plane.

It is convenient to consider also the correlation function of two Euclidean Wilson loops $\tilde{\mathcal{W}}_1$ and $\tilde{\mathcal{W}}_2$ running along two rectangular paths $\tilde{\mathcal{C}}_1$ and $\tilde{\mathcal{C}}_2$ which follow the following straight-line trajectories:

$$\begin{aligned}\tilde{\mathcal{C}}_1 &\rightarrow X_{E\mu}^{(\pm 1)}(\tau) = z_{E\mu} + \frac{p_{1E\mu}}{m}\tau \pm \frac{R_{1E\mu}}{2}, \\ \tilde{\mathcal{C}}_2 &\rightarrow X_{E\mu}^{(\pm 2)}(\tau) = \frac{p_{2E\mu}}{m}\tau \pm \frac{R_{2E\mu}}{2},\end{aligned}\tag{17}$$

and are closed by straight-line paths at proper times $\tau = \pm T$. Here $R_{1E} = (0, \vec{R}_{1\perp}, 0)$, $R_{2E} = (0, \vec{R}_{2\perp}, 0)$ and $z_E = (0, \vec{z}_\perp, 0)$. Moreover, in the Euclidean theory we *choose* the four-vectors p_{1E} and p_{2E} to be:

$$\begin{aligned}p_{1E} &= m(\sin \frac{\theta}{2}, 0, 0, \cos \frac{\theta}{2}), \\ p_{2E} &= m(-\sin \frac{\theta}{2}, 0, 0, \cos \frac{\theta}{2}),\end{aligned}\tag{18}$$

$\theta \in]0, \pi[$ being the angle formed by the two trajectories (+1) and (+2) in Euclidean four-space. Let us introduce the following notations for the normalized correlators $\langle \mathcal{W}_1 \mathcal{W}_2 \rangle / \langle \mathcal{W}_1 \rangle \langle \mathcal{W}_2 \rangle$ in the Minkowskian and in the Euclidean theory, in the presence of a *finite* IR cutoff T :

$$\begin{aligned}\mathcal{G}_M(\chi; T; \vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp}) &\equiv \frac{\langle \mathcal{W}_1^{(T)} \mathcal{W}_2^{(T)} \rangle}{\langle \mathcal{W}_1^{(T)} \rangle \langle \mathcal{W}_2^{(T)} \rangle}, \\ \mathcal{G}_E(\theta; T; \vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp}) &\equiv \frac{\langle \tilde{\mathcal{W}}_1^{(T)} \tilde{\mathcal{W}}_2^{(T)} \rangle_E}{\langle \tilde{\mathcal{W}}_1^{(T)} \rangle_E \langle \tilde{\mathcal{W}}_2^{(T)} \rangle_E}.\end{aligned}\tag{19}$$

As already stated in Ref. ⁵, the two quantities in Eq. (19) (with $\chi > 0$ and $0 < \theta < \pi$) are expected to be connected by the same analytic continuation in the angular variables and in the IR cutoff which was already derived in the case of Wilson lines ^{5,6,7}, i.e.:

$$\begin{aligned}\mathcal{G}_E(\theta; T; \vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp}) &= \mathcal{G}_M(\chi \rightarrow i\theta; T \rightarrow -iT; \vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp}), \\ \mathcal{G}_M(\chi; T; \vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp}) &= \mathcal{G}_E(\theta \rightarrow -i\chi; T \rightarrow iT; \vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp}).\end{aligned}\tag{20}$$

Indeed it can be proved¹¹, simply by adapting step by step the proof derived in Ref. ⁵ from the case of Wilson lines to the case of Wilson loops, that the analytic continuation (20) is an *exact* result, i.e., not restricted to some order in perturbation theory or to some other approximation, and is valid both for the Abelian and the non-Abelian case.

By using the analytic continuation (20), one can also derive the following *crossing-symmetry* relation:

$$\begin{aligned}\mathcal{G}_M(i\pi - \chi; T; \vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp}) &= \mathcal{G}_M(\chi; T; \vec{z}_\perp, \vec{R}_{1\perp}, -\vec{R}_{2\perp}) \\ &= \mathcal{G}_M(\chi; T; \vec{z}_\perp, -\vec{R}_{1\perp}, \vec{R}_{2\perp}).\end{aligned}\quad (21)$$

As we have said above, the loop-loop correlation functions (19), both in the Minkowskian and in the Euclidean theory, are expected to be IR-*finite* quantities, i.e., to have finite limits when $T \rightarrow \infty$, differently from what happens in the case of Wilson lines. One can then define the following loop-loop correlation functions with the IR cutoff removed:

$$\begin{aligned}\mathcal{C}_M(\chi; \vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp}) &\equiv \lim_{T \rightarrow \infty} [\mathcal{G}_M(\chi; T; \vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp}) - 1], \\ \mathcal{C}_E(\theta; \vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp}) &\equiv \lim_{T \rightarrow \infty} [\mathcal{G}_E(\theta; T; \vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp}) - 1].\end{aligned}\quad (22)$$

As a pedagogic example to illustrate these considerations, we shall consider the simple case of QED, in the so-called *quenched* approximation, where vacuum polarization effects, arising from the presence of loops of dynamical fermions, are neglected. In this approximation, the calculation of the normalized correlators (19) can be performed exactly (i.e., without further approximations) both in Minkowskian and in Euclidean theory and one finds that¹¹ i) the two quantities $\mathcal{G}_M$ and $\mathcal{G}_E$ are indeed connected by the analytic continuation (20), and ii) the two quantities are finite in the limit when the IR cutoff T goes to infinity:

$$\mathcal{C}_M(\chi; \vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp}) = \exp \left[-i4e^2 \coth \chi \, t(\vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp}) \right] - 1, \quad (23)$$

$$\mathcal{C}_E(\theta; \vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp}) = \exp \left[-4e^2 \cot \theta \, t(\vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp}) \right] - 1, \quad (24)$$

where

$$t(\vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp}) \equiv \frac{1}{8\pi} \ln \left(\frac{|\vec{z}_\perp + \frac{\vec{R}_{1\perp}}{2} + \frac{\vec{R}_{2\perp}}{2}| |\vec{z}_\perp - \frac{\vec{R}_{1\perp}}{2} - \frac{\vec{R}_{2\perp}}{2}|}{|\vec{z}_\perp + \frac{\vec{R}_{1\perp}}{2} - \frac{\vec{R}_{2\perp}}{2}| |\vec{z}_\perp - \frac{\vec{R}_{1\perp}}{2} + \frac{\vec{R}_{2\perp}}{2}|} \right). \quad (25)$$

As shown in Ref. ¹¹, the results (23) and (24) can be used to derive the corresponding results in the case of a non-Abelian gauge theory with N_c colours, up to the order $\mathcal{O}(g^4)$ in perturbation theory (see also Refs. ^{12,13}):

$$\mathcal{C}_M(\chi; \vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp})|_{g^4} = -2g^4 \left(\frac{N_c^2 - 1}{N_c^2} \right) \coth^2 \chi [t(\vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp})]^2, \quad (26)$$

$$\mathcal{C}_E(\theta; \vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp})|_{g^4} = 2g^4 \left(\frac{N_c^2 - 1}{N_c^2} \right) \cot^2 \theta [t(\vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp})]^2. \quad (27)$$

We stress the fact that both the Minkowskian quantities (23) and (26) and the Euclidean quantities (24) and (27) are IR finite, differently from the corresponding quantities constructed with Wilson lines, which were evaluated in Ref. ⁶ (see also Ref. ²).

It is also important to notice that the two quantities (23) and (24), as well as the two quantities (26) and (27), obtained *after* the removal of the IR cutoff ($T \rightarrow \infty$), are still connected by the usual analytic continuation in the angular variables only:

$$\begin{aligned}\mathcal{C}_E(\theta; \vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp}) &= \mathcal{C}_M(\chi \rightarrow i\theta; \vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp}), \\ \mathcal{C}_M(\chi; \vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp}) &= \mathcal{C}_E(\theta \rightarrow -i\chi; \vec{z}_\perp, \vec{R}_{1\perp}, \vec{R}_{2\perp}).\end{aligned}\quad (28)$$

[Moreover, the expressions (23) and (26) trivially satisfy the crossing–symmetry relation (21).] This is a highly non–trivial result, whose general validity is discussed in Ref. ¹¹. (Indeed, the validity of the relation (28) has been also recently verified in Ref. ¹³ by an explicit calculation up to the order $\mathcal{O}(g^6)$ in perturbation theory.)

As said in Ref. ¹¹, if $\mathcal{G}_M$ and $\mathcal{G}_E$, considered as functions of the *complex* variable T , have in $T = \infty$ an “eliminable *isolated* singular point” [i.e., they are analytic functions of T in the *complex* region $|T| > R$, for some $R \in \mathbb{R}^+$, and the *finite* limits (22) exist when letting the *complex* variable $T \rightarrow \infty$], then, of course, the analytic continuation (28) immediately derives from Eq. (20) (with $|T| > R$), when letting $T \rightarrow +\infty$. (For example, if $\mathcal{G}_M$ and $\mathcal{G}_E$ are analytic functions of T in the *complex* region $|T| > R$, for some $R \in \mathbb{R}^+$, and they are bounded at large T , i.e., $\exists B_{M,E} \in \mathbb{R}^+$ such that $|\mathcal{G}_{M,E}(T)| < B_{M,E}$ for $|T| > R$, then $T = \infty$ is an “eliminable singular point” for both of them.) But the same result (28) can also be derived under different conditions. For example, let us assume that $\mathcal{G}_E$ is a bounded analytic function of T in the sector $0 \leq \arg T \leq \frac{\pi}{2}$, with finite limits along the two straight lines on the border of the sector: $\mathcal{G}_E \rightarrow G_{E1}$, for $(\text{Re}T \rightarrow +\infty, \text{Im}T = 0)$, and $\mathcal{G}_E \rightarrow G_{E2}$, for $(\text{Re}T = 0, \text{Im}T \rightarrow +\infty)$. And, similarly, let us assume that $\mathcal{G}_M$ is a bounded analytic function of T in the sector $-\frac{\pi}{2} \leq \arg T \leq 0$, with finite limits along the two straight lines on the border of the sector: $\mathcal{G}_M \rightarrow G_{M1}$, for $(\text{Re}T \rightarrow +\infty, \text{Im}T = 0)$, and $\mathcal{G}_M \rightarrow G_{M2}$, for $(\text{Re}T = 0, \text{Im}T \rightarrow -\infty)$. We can then apply the “Phragmén–Lindelöf theorem” (see, e.g., Theorem 5.64 in Ref. ¹⁴) to state that $G_{E2} = G_{E1}$ and $G_{M2} = G_{M1}$. Therefore, also in this case, the analytic continuation (28) immediately derives from Eq. (20) when $T \rightarrow \infty$.

The relation (28) has been extensively used in the literature ^{12,15,16,17,18} in order to address, from a non–perturbative point of view, the still unsolved problem of the asymptotic s –dependence of hadron–hadron elastic scattering amplitudes and total cross sections. (See, e.g., Ref. ¹⁹ and references therein for a recent review of the problem. It has been also recently proved in Ref. ¹³, by an explicit perturbative calculation, that the loop–loop scattering amplitude approaches, at sufficiently high energy, the BFKL–*pomeron* behaviour ²⁰.)

An independent non–perturbative approach would be surely welcome and could be provided by a direct lattice calculation of the loop–loop Euclidean correlation functions. This would surely result in a considerable progress along this line of research.

References

1. O. Nachtmann, Ann. Phys. **209** (1991) 436.
2. E. Meggiolaro, Phys. Rev. D **53** (1996) 3835.
3. E. Meggiolaro, Nucl. Phys. B **602** (2001) 261.
4. H. Verlinde and E. Verlinde, hep–th/9302104.
5. E. Meggiolaro, Nucl. Phys. B **625** (2002) 312.
6. E. Meggiolaro, Z. Phys. C **76** (1997) 523.
7. E. Meggiolaro, Eur. Phys. J. C **4** (1998) 101.
8. O. Nachtmann, in *Perturbative and Nonperturbative aspects of Quantum Field Theory*, edited by H. Latal and W. Schweiger (Springer–Verlag, Berlin, Heidelberg, 1997).
9. H.G. Dosch, E. Ferreira and A. Krämer, Phys. Rev. D **50** (1994) 1992.
10. E.R. Berger and O. Nachtmann, Eur. Phys. J. C **7** (1999) 459.
11. E. Meggiolaro, Nucl. Phys. B **707** (2005) 199.
12. A.I. Shoshi, F.D. Steffen, H.G. Dosch and H.J. Pirner, Phys. Rev. D **68** (2003) 074004.
13. A. Babansky and I. Balitsky, Phys. Rev. D **67** (2003) 054026.
14. E.C. Titchmarsh, *The theory of functions*, second edition (Cambridge University Press, London, 1939).

15. R.A. Janik and R. Peschanski, Nucl. Phys. B **586** (2000) 163.
16. R.A. Janik, Phys. Lett. B **500** (2001) 118.
17. E. Shuryak and I. Zahed, Phys. Rev. D **62** (2000) 085014.
18. A.E. Dorokhov and I.O. Cherednikov, Ann. Phys. **314** (2004) 321.
19. S. Donnachie, G. Dosch, P. Landshoff and O. Nachtmann, *Pomeron Physics and QCD* (Cambridge University Press, Cambridge, 2002).
20. V.S. Fadin, E.A. Kuraev and L.N. Lipatov, Phys. Lett. B **60** (1975) 50;
I.I. Balitsky and L.N. Lipatov, Sov. Journ. Nucl. Phys. **28** (1978) 822.